

SCATTERING-TYPE NEAR-FIELD MICROSCOPY: FROM NANOSCALE INFRARED MATERIAL RECOGNITION TO SUPERLENS STUDIES

R. Hillenbrand

Nano-Photonics Group, Max-Planck-Institut für Biochemie, 82152 Martinsried, Germany
hillenbr@biochem.mpg.de

The development of novel nanoelectronic and photonic structures requires ultrahigh-resolution optical microscopy for characterization and mapping of local material properties and nanoscale confined light fields. I will demonstrate such an optical microscopy technique providing a spatial resolution of about 10-20 nm independent of the wavelength. It is based on elastic light scattering from the probing tip of an atomic force microscope (scattering-type near-field optical microscopy, s-SNOM [1]). Besides a short introduction of the technique, I will demonstrate some s-SNOM applications such as mid-infrared mapping of material [2,3] and doping contrasts in semiconductor devices [4], single nanoparticle mapping [5,6] and near-field characterization of a SiC superlens [7].




Fig. 1: Nanometer-scale resolved s-SNOM images of a cross section of a MOS-transistor drain contact recorded at a wavelength of about 10.8 μm . Pseudoheterodyne infrared detection yields amplitude and phase images showing distinct material and doping contrasts simultaneously to topography.

References:

- [1] F. Keilmann and R. Hillenbrand, *Philos. Trans. R. Soc. London, Ser. A* **362**, 739 (2004).
- [2] R. Hillenbrand, T. Taubner, and F. Keilmann, *Nature* **418**, 159 (2002).
- [3] N. Ocelic and R. Hillenbrand, *Nat. Mater.* **3**, 606 (2004).
- [4] A. Huber, D. Kazantsev, F. Keilmann, J. Wittborn, and R. Hillenbrand, *Adv. Mater.*, published online (2007)
- [5] A. Cvitkovic, N. Ocelic, J. Aizpurua, R. Guckenberger, and R. Hillenbrand, *Phys. Rev. Lett.* **97**, 060801 (2006).
- [6] M. Brehm, T. Taubner, R. Hillenbrand, and F. Keilmann, *Nano Lett.* **6**, 1307 (2006).
- [7] T. Taubner, D. Korobkin, Y.A. Urzhumov, G. Shvets, and R. Hillenbrand *Science*, **313**, 1595 (2006).